

Learning together to be  
the best we can be

Headteacher - Nichola Irving



# Supporting Pupils with Medical Conditions

## Safeguarding Statement

**Everyone at Horton Grange shares an objective to help keep children and young people safe by contributing to:**

- Providing a safe environment for children and young people to learn in school and;
- Identifying children and young people who are suffering or likely to suffer significant harm, and taking appropriate action with the aim of making sure they are kept safe both at home and in school.

Date adopted: July 2014  
Reviewed date: September 2019  
Next review date: September 2020

Lifelong learning starts here...

## **Contents**

---

Contents .....	1
Foreword.....	2
Roles and Responsibilities .....	2
General Procedures .....	5
Refusal or Forgetting to Take Medication.....	6
Non-prescribed Medication .....	6
Individual Healthcare Plan .....	7
Practical Advice for Common Conditions .....	8
Anaphylaxis .....	9
Asthma .....	11
Diabetes.....	19
Epilepsy .....	22
Unacceptable Practice .....	26
Complaints.....	27
Administration of Medication to Pupils.....	28

## **Appendices**

Agreement between Parents and School (Appendix 1) .....	28
Parental Request for Child to Carry and Self-administer Medicine (Appendix 2) .....	30
Healthcare Plan for a Pupil with Medical Needs (Appendix 3) .....	31
Individual Staff Training Record – Administration of Medication (Appendix 4) .....	34
Record of Medication Administered in School (Appendix 5) .....	35
Parental Consent: Use of Emergency Salbutamol Inhaler Appendix 6 .....	36
Specimen letter to inform parents that the emergency salbutamol inhaler was used (Appendix 7) .....	37
Emergency Action in the Event of an Asthma Attack (Appendix 8).....	38
Emergency Action: Epilepsy - First Aid for all Seizures (Appendix 9) .....	39
Emergency Action: First Aid for Children Known to Have Epilepsy and Prescribed Rectal Diazepam (Appendix 10).....	40
Individual Care Plan for the Administration of Rectal Diazepam (Appendix 11).....	41
Record of Use of Rectal Diazepam (Appendix 12) .....	44

## **Foreword**

---

The County Council's Corporate Health and Safety Team has prepared this guidance in consultation with paediatricians from Northumbria Healthcare NHS Foundation Trust. It chiefly refers to essential prescribed medication. The contents fully complement information contained in the Department for Education's (DfE's) document entitled '[Supporting pupils at school with medical conditions](#)' which was published in April 2014. It also supersedes the document 'Policy on Supporting Children with Medical Needs' which the Council published in July 1999. Schools should dispose of all previous copies of the latter guidance.

## **Roles and Responsibilities**

---

### **Responsibility of Parents**

In modern terminology the term 'parent' is understood to mean not just a parent but anyone who has parental responsibility for the care of a child. Parents have the principal responsibility for the administration of medication to their children, who have the right to be educated with their peers, regardless of any short or long-term needs for medication whilst at school.

It is preferable that medication be given at home whenever possible. If prescribed medicines are to be taken three or more times per day, parents should ask the prescribing doctor if the administration of the medication can occur outside normal school hours. Non-prescription medication (such as cough medicines) should not be administered in school. However, in certain circumstances analgesics can be given (see page 4).

The parents have the principal duty to inform the school of their children's medical conditions and to make a request for the Head or Medication Coordinator to make arrangements for medication to be administered in school. This can occur if the child:

- has been newly diagnosed
- is due to return after a long absence and has a chronic illness or long-term complaints, such as asthma, diabetes, epilepsy or another condition
- is recovering from a short-term illness and is well enough to return to school whilst still receiving a course of antibiotics or other medication.
- has needs that have changed
- is due to attend a new school

## **Responsibility of Health Care Professionals**

In situations where the condition requires a detailed individual healthcare plan or specific specialist training is required for school staff this will often require direct input from Healthcare Professionals with clinical responsibility for the child. Examples include community or specialist nurses and, in the case of children with mobility needs, occupational therapists or physiotherapists.

Often the specific details in an individual healthcare plan can only be provided by professionals who have access to the confidential notes that the Consultants and other healthcare professionals working with the child in question have prepared.

The roles of the Occupational Therapists and physiotherapists are clear; schools should continue to contact them at Wansbeck General Hospital. The main switchboard number is 0344 811 8111.

The School Nursing Team is able to provide training on anaphylaxis and can provide a 'signposting role' should schools have difficulty accessing professional medical assistance or if there is uncertainty about which consultant to contact.

The specialist nurses employed by the Northumbria Trust are hospital-based and work directly with the relevant consultants. They provide training for diabetes and epilepsy and can offer the necessary assistance with healthcare plans for these conditions.

## **Responsibility of School Staff**

Each request for medicine to be administered to a pupil in school should be considered on its merits. The Head should give consideration to the best interests of the pupil and the implications for the school.

It is generally accepted that school staff may administer prescribed medication whilst acting in loco parentis. However, it is important to note that this does not imply that there is a duty upon these workers to administer medication and the following should be taken into account:

No member of staff should be compelled to administer medication to a pupil

No medication can be administered in school without the agreement of Mrs Irving Headteacher or Mrs Claire Jameson SENDCO/Medication Coordinator

The Head and governors must nominate a member of staff to assume the role of Medication Coordinator, who will have overall responsibility for the implementation of this policy. In addition to this, if schools have an emergency asthma kit, the Department of Health recommends at least two volunteers should have the responsibility of ensuring this kit is maintained correctly.

Separate registers must be compiled defining those pupils who are permitted to use emergency inhalers and emergency AAI.

There are named personnel who administer medicine :

In EYFS this is Kerry Graham, Wendy Cochrane and David Hansen

In KS1 this is Karen Adam and Sarah Foy

In KS2 this is Pauline Alderson, Joanne Dyson Julie Slater

The volunteers should receive the appropriate guidance and training (where necessary)

Parents requesting administration of medication form for their children should be referred to the school's webpage or they can collect one from Reception. They should be asked to complete Part 1 of the form 'Administration of Medication to Pupils – Agreement between Parents and School'. Completion of this form safeguards staff by allowing only prescribed medication to be administered

School staff may consult with the prescriber to ascertain whether medication can be given outside of school hours.

## **Day Trips, Residential Visits and Sporting Activities**

School staff should be aware of how a child's medical condition will impact on their participation in such events, however, there should be enough flexibility for all children to participate according to their abilities and with any reasonable adjustments. The only exception would be in cases where evidence from a clinician, such as a GP or specialist consultant, states that participation is not possible.

To determine the reasonable adjustments required, the school should complete a personalised risk assessment. This will ensure that planning arrangements take account of any steps needed to ensure that pupils with medical needs are included. The risk assessment will require consultation with parents and pupils and advice from the relevant healthcare professionals to ensure that pupils can participate safely. In certain situations it may also be appropriate to consult the intended activity providers.

The specific arrangements for a child attending an off-site visit should be recorded with the relevant visit specific information and logged on the EVOLVE system, together with supporting information, such as the risk assessment. These arrangements enable the child to participate fully in such activities and ensure social inclusion, as recommended by the Outdoor Education Advisory Board's National guidance 3.2e 'Inclusion'.

In general, the arrangements detailed in this policy for the safe storage and administration of medication should also apply when taken off site. Medication should be stored in a locked bag or box and held by a designated member of staff who is trained to administer it. Associated documentation, such as the personalised risk assessment, Healthcare plan and Medical Administration Record (MAR) (for Special Schools) should also be available and securely stored with the medication or on an encrypted digital device (which does not rely on a data signal to access it).

The EVOLVE system has a facility which allows a visit form along with all associated attachments to be downloaded electronically to a Compressed (zipped) folder to a digital device. This may assist those schools wishing to store documents on an encrypted digital platform.

Emergency medication, such as inhalers, AAIs etc, should be readily available and not locked away. A record of any medication administered should be kept.

## **Liability and Indemnity**

Members of staff administering medication in accordance with appropriate training or the details supplied by the parent may rest assured that they are indemnified under the conditions of the existing insurance policies. In such circumstances, any liabilities rest with the insured party.

## General Procedures

---

- If medication cannot be given outside of school hours, parents should fill in the aforementioned request form (Appendix 1) giving the dose to be taken, the method of administration, the time and frequency of administration, other treatment, any special precautions and signed consent.
- The parent (not the pupil) should bring all essential medication to school. It should be delivered personally to the Office staff who will pass it to the Medication coordinators. Only the smallest practicable amount should be kept in school.
- All medication taken in school must be kept in a clearly labelled pharmacy bottle, preferably with a child safety top, which must have the owner's name, photograph, the contents and the dosage to be administered.
- Whilst medication is in school it should be kept in a locked cupboard or fridge (if so required), preferably in the staffroom or main office. In the event of an emergency it should be readily accessible to the named volunteer or young person, when required. The exceptions to this are inhalers, adrenaline auto-injectors and insulin. These medications should be carried by the child or may be kept in the classroom, depending on the child's age and developing independence.
- In the case of epilepsy medication and diabetes monitoring kits they should be carried in a bum bag or rucksack by the adult with the child. They should be worn at all times. Epilepsy medication will be in a child-proof container.
- Medication to be taken orally should be supplied with an individual measuring spoon or syringe. Eye drops and ear drops should be supplied with a dropper. A dropper or spoon must only be used to administer medicine to the owner of that implement.
- When medication is given, the name of the drug, the dose, the mode of administration, the time that treatment is required to be given and date of expiry should be checked. A written record should be kept of the time it was given and by whom to avoid more than one person ever giving more than the recommended dose. This should be kept with the parental consent form. See form in Appendix 1.
- Where any change of medication or dosage occurs, clear written instructions from the parent should be provided. If a pupil brings any medication to school for which consent has not been given, school staff can refuse to administer it. In such circumstances the Head or Medical Coordinator should contact the parent as soon as possible.
- Renewal of medication which has passed its expiry date is the responsibility of the parent. Nevertheless, schools should have robust procedures in place to ensure that out of date medication is not administered in error. If parents are unable to collect expired medication then staff should take it to the local pharmacy so that it can be disposed of safely. The medication must not be disposed of in any other way.
- In all cases where, following the administration of medication, there are concerns regarding the reaction of the pupil, medical advice should be sought immediately and the parents informed.
- On transition to a new school, parents will be handed back any medication in school. All records will be handed on to the new school.

If members of staff are in doubt about any of the above procedures they should check with the parents or a health professional before taking further action.

## **Refusal or Forgetting to Take Medication**

---

If pupils refuse medication or forget to take it, the school should inform the child's parent as a matter of urgency. If necessary, the school should call the emergency services.

## **Non-prescribed Medication**

---

As a rule, non-prescribed (over-the-counter) medication, such as cough medicines, should not be administered. However, there is one notable exception to this rule, namely, paracetamol.

At the discretion of the Head, paracetamol can be issued, provided the practice is strictly controlled by adopting the same standards as for prescribed medication. Once again, a formal agreement should be made between the school and the parents (see Appendix 1 below).

The Head should authorise specific members of staff to dispense the tablets. In order to monitor and prevent the danger of any individuals overdosing on the medication the nominated member of staff should keep a record of when it was issued, giving such information as name of the pupil and the time and the dose which was administered (see Appendix 5). Before administering the medication members of staff should always ask the child whether any side effects or allergic reactions have been experienced.

The paracetamol must be kept securely under lock and key and dispensed with care since over-dosage is dangerous. Paracetamol must not be kept in first-aid boxes.

In the main, children of First or Primary School age should only be given paracetamol in exceptional circumstances under the supervision of their GP. However, in certain circumstances, such as if a child has a raised temperature, the Head can authorise the use of paracetamol via a formal agreement with the parent, as described above.

On no account should aspirin or preparations that contain aspirin be given to pupils unless a doctor has prescribed such medication.

## **Individual Healthcare Plan**

---

This section of the policy covers the role of individual healthcare plans in supporting pupils at school who have long-term, severe or complex medical conditions. The new statutory guidance imposes a requirement to identify the member of staff who is responsible for the development of these plans. In school this is Claire **Jameson**.

The governing body should ensure that there are robust school arrangements to:

- establish the need for a plan
- ensure that plans are adequate
- review plans at least annually or earlier if evidence indicating that the child's needs have changed is brought to its attention.

Healthcare plans should be developed with the child's best interests in mind and the school should ensure that it assesses and manages risks to the child's education, health and social well-being and minimises disruption.

Personalised risk assessments, moving and handling risk assessments, emergency procedures and other such documents should be used to supplement the individual healthcare plan, as appropriate.

The schools healthcare plan is given in Appendix 3. To ensure compliance with the new statutory guidance the following issues have been taken into account:

- the medical condition, its triggers, signs, symptoms and treatments
- the pupil's resulting needs, including medication (with details of dose, side-effects and storage arrangements) and other treatments, time, facilities, equipment, testing, access to food and drink where this is used to manage his/her condition, dietary requirements and environmental issues such as crowded corridors, travel time between lessons
- specific support for the pupil's educational, social and emotional needs – for example, how absences will be managed, requirements for extra time to complete exams, use of rest
- periods or additional support in catching up with lessons, counselling sessions
- the level of support needed, (some children will be able to take responsibility for their own health needs), including in emergencies. If a child is self-managing medication to be clearly stated with appropriate periods of monitoring
- who will provide this support, their training needs, expectations of their role and confirmation of their proficiency to provide support for the child's medical condition from a healthcare professional, together with an indication of the arrangements for cover that will be available when those supporting are unavailable
- who in the school needs to be aware of the child's condition and the support required
- the need to establish arrangements which enable written permission from parents and the Head to be drawn up, thus authorising a member of staff to administer medication or allowing the pupil to self-administer during school hours
- the designated individuals to be entrusted with information about the child's condition where the parent or child has raised confidentiality issues
- what to do in an emergency, including whom to contact, and contingency arrangements. Some children may have an emergency healthcare plan prepared by their lead clinician that could be used to inform development of their individual healthcare plan.
- the separate arrangements or procedures required for school trips, educational visits or other extra-curricular activities. In practice, these should be logged on the EVOLVE system, together with supporting information, such as personalised risk assessments. These arrangements enable the child to participate fully in such activities and ensure social inclusion, as recommended by the Outdoor Education Advisory Board's National guidance 3.2e 'Inclusion'.

## Practical Advice for Common Conditions

---

A small number of children need medication to be given by injection, auto-injectors or other routes. The most appropriate arrangements for managing these situations effectively are best determined by agreement between the school, parent, school nurse (where there is one) and the doctor who prescribed the medication.

Members of staff willing to administer medication should be made fully aware of the procedures and should receive appropriate training from competent healthcare staff. More information on training requirements is given below in the sections of this policy covering common medical conditions. The majority of parents will be aware of the contact details for their child's specialist nurse. Schools should contact them directly in the first instance. The School Nursing Team can be contacted for advice and is able to direct inquirers to other health agencies, where necessary. An individual healthcare plan for each pupil with a medical need must be completed and conform to the procedures described on pages 5 and 6. Information in the appendices should prove helpful.

The medical conditions in children that most commonly cause concern in schools are asthma, epilepsy, diabetes and anaphylaxis. Essential information about these conditions is provided within this policy. More detailed information can be obtained from the following organisations:

- [Asthma UK](#)
- [Epilepsy Society](#) (formerly The National Society for Epilepsy)
- [Epilepsy Action](#) (formerly the British Epilepsy Association)
- [Diabetes UK](#)
- [Anaphylaxis Campaign](#)
- [National Electronic Library for Medicines \(NHS\)](#)
- [Resuscitation Council \(UK\)](#)

## Anaphylaxis

---

### What is Anaphylaxis?

Anaphylaxis is an extreme allergic reaction that occurs rarely in people who have an extreme sensitivity to a particular substance known as an allergen. It can affect the whole body, including the airways and circulation. Often it occurs within minutes of exposure to the allergen, though sometimes it does not arise until many hours later. **Anaphylaxis is potentially life threatening - delays in giving adrenaline are a common finding in fatal reactions.**

### What Causes it?

Common causes of anaphylaxis include:

- Edible triggers, such as peanuts, tree nuts, fish, shellfish, dairy products and eggs
- Other triggers, such as natural latex, the venom of stinging insects (for example wasps, bees and hornets) penicillin and any other drugs or injections

Anaphylactic shock is the most severe form of allergic reaction. This occurs when the blood pressure falls dramatically and the patient loses consciousness.

## What are the Signs of the Condition?

Common signs of anaphylaxis in children include:

- swelling in the throat, which can restrict the air supply thus causing breathing difficulties.
- severe asthma
- dizziness
- itchy skin, generalised flushing of the skin, tingling or itching in the mouth or hives anywhere on the body
- swelling of the lips, hands and feet
- abdominal cramps, nausea and vomiting.

## What is the Treatment for the Condition?

The treatment for a severe allergic reaction is an injection of adrenaline (also known as epinephrine) into the muscle of the upper outer thigh via a pre-loaded injection device, such as an epiPen, anapen or jext. An injection should be given as soon as a reaction is suspected.

**Anaphylaxis is potentially life threatening - delays in giving adrenaline are a common finding in fatal reactions** Anaphylaxis should always be regarded as a medical emergency which requires that an ambulance be called immediately. **Always make it clear to the emergency operator that the child is showing symptoms of anaphylaxis so that they can prioritise the call.**

## What Arrangements are in Place at our School?

Medication is provided by the parents at the beginning of every school year or when needed. It is handed back at the end of the school year or on the expiration date whichever is sooner.

Children who have known allergies have a completed Health Plan. These plans are updated annually. All members of staff who work with that child are aware of the plan. Staff have regular training on the administration of medicines.

All medication for its treatment is kept unlocked in the classroom in a place which is easily accessible.

An ambulance will be called as soon as possible.

## **Healthcare Plan**

Anaphylaxis is manageable. With sound precautionary measures, the development of a suitable healthcare plan and support from members of staff, school life may continue as normal for all concerned.

It is important that appropriate local procedures for the use of adrenaline auto-injectors, should include the following

- awareness among all members of staff that the child has this particular medical condition
- awareness of the symptoms associated with anaphylactic shock
- knowledge of the type of injector to be used
- labelling of injectors for the child concerned, for example adrenaline, anti-histamine
- knowledge of the locations where the injector is stored, preferably in an easily accessible place such as a medication box
- the provision of appropriate instruction and training to nominated members of staff
- familiarity with the names of those trained to administer treatment
- an understanding of the need to keep records of the dates of issue knowledge of emergency contacts

This type of information should be suitably displayed in the areas where the medication is to be kept. This information should include the name of the child and a photograph. Care must be given to ensure confidentiality. The information should be accessible but not publicly displayed. Information should accompany the medication on school trips. The arrangements for swimming and other sporting activities should also be considered.

Collectively, it is for the Head, the child's parents and the medical staff involved to decide how many adrenaline devices the school should hold, and where they should be stored.

Where children are deemed sufficiently responsible for carrying their own emergency treatment with them, it is nevertheless important that a spare set should always be kept safely on site. This should be accessible to all staff and stored in a secure place. In large schools or split sites, it is often quicker for staff to use an injector that is with the child rather than taking time to collect one from a central location. In an emergency situation it is important to avoid any delay.

## **Spare Emergency AAI**

**It is important that school staff read the content of the DoH document  
Guidance on the use of adrenaline auto-injectors in Schools dated September 2017.**

## **Supply**

The spare AAI can be administered to a pupil whose own prescribed AAI cannot be administered correctly without delay or is not available; it does not replace a child's own prescribed AAI. Staff are also able to administer this to those children who have not been prescribed an AAI, but only in cases where both written medical authorisation and written parental consent for its use has been provided. In the event of a possible severe allergic reaction in a pupil who does not meet these criteria, emergency services (999) should be contacted and advice sought from them as to whether administration of the spare AAI is appropriate.

## **Protocols for Emergency AAI**

- Spare AAI's will be kept in corridor first aid cupboards.
- School will have a register of pupils who have been prescribed an AAI
- written consent from the pupil's parent/legal guardian for use of the spare AAI(s), as part of a pupil's individual healthcare plan.
- ensuring that any spare AAI is used only on pupils where both medical authorisation and written parental consent has been provided.
- appropriate support and training for staff in the use of the AAI in line with the schools wider policy on supporting pupils with medical conditions.
- arrangements for purchase, storage and disposal of spare AAIs.
- Nicola Irving and Claire Jameson will be responsible for the implementation and monitoring of the protocols.
- School will keep a record of use of any AAI(s), as required by Supporting Pupils policy and inform parents or carers that their pupil has been administered an AAI and whether this was the school's spare AAI or the pupil's own device.
- All staff will know where emergency AAI's are kept (In area First Aid boxes) and be able to access them.

## **Food Management**

Day-to-day policy measures are needed for food management; awareness of the child's needs in relation to the menu, individual meal requirements and snacks in school. When catering staff are employed by a separate organisation, it is important to ensure that the Unit Manager is fully aware of the child's particular requirements. A 'kitchen code of practice' should be put in place.

Parents often ask for the Head to exclude from the premises the food to which their child is allergic. This is not always feasible, although appropriate steps to minimise any risks to allergic children should be taken.

## Training

Staff will:

- be trained to recognise the range of signs and symptoms of an allergic reaction;
- understand the rapidity with which anaphylaxis can progress to a life-threatening reaction, and that anaphylaxis may occur with prior mild (e.g. skin) symptoms;
- appreciate the need to administer adrenaline without delay as soon as anaphylaxis occurs, before the patient might reach a state of collapse (after which it may be too late for the adrenaline to be effective);
- be aware of the anaphylaxis policy;
- be aware of how to check if a pupil is on the register;
- be aware of how to access the AAI.

## Asthma

---

### What is Asthma?

People with asthma have airways which narrow as a reaction to various triggers. The narrowing or obstruction of the airways causes breathing difficulties.

### What Causes it?

There are many things that can trigger an asthma attack. Common examples include:

- viral infections
- house dust mites
- pollen
- smoke
- fur
- feathers
- pollution
- laughter
- excitement
- stress

### What are the Signs of the Condition?

The most common symptoms of an asthma attack include:

- coughing
- wheezing
- difficulty breathing
- nasal flaring
- a tight feeling in the chest (younger children may express this as 'tummy ache' or feeling like someone is sitting on their chest)
- Inability to talk or complete sentences (some children will go very quiet).

## **What is the Treatment for the Condition?**

The main types of medicines used to treat asthma are discussed briefly below:

### **Relievers**

Usually it is a reliever that a child will need during the school day. Relievers (usually blue inhalers) are medicines that are taken immediately to relieve the symptoms of asthma during an attack. They quickly relax the muscles surrounding the narrowed airways thus allowing them to open wider, making it easier for the child to breathe. They are sometimes taken before exercise.

### **Preventers**

Preventer inhalers can be brown, red or orange in colour and can sometimes be in the form of tablets. Preventers are usually used out of school hours and it is rare for them to be needed during the school day.

Preventers protect the lining of the airways, help to calm the swelling and stop the tubes in the lungs from being so sensitive.

### **Spacers**

Both kinds of inhalers are often used in combination with spacers which help deliver medicine to the lungs more effectively. Where prescribed, the spacer should be individually labelled with the child's name and kept with the inhaler.

### **Nebulisers**

A nebuliser is a machine that creates a mist of medicine that is then breathed through a mask or mouthpiece. They are becoming increasingly less common. Pupils with asthma should not normally need to use a nebuliser in school. However, if they do have to use one then members of school staff will need to receive appropriate training from a healthcare professional.

### **Training**

Since emergency treatments vary in each case, the parents will often be best placed to inform schools of the child's treatment regime. There may be a specialist nurse from the local NHS Trust who can deliver training and will have access to the medical advice that has informed the healthcare plan.

Children with asthma will often be looked after solely by their GP or Asthma Nurse. Although the GP would be unable to provide training it is likely that they will provide the information that would help school staff to complete the healthcare plans. Children with complex conditions may have access to a specialist nurse with expert knowledge in oncology, nephrology, gastroenterology, urology or cystic fibrosis, who may be able to assist.

## Designated Members of Staff

Designated members of staff should be trained in:

- recognising asthma attacks (and distinguishing them from other conditions with similar symptoms)
- responding appropriately to a request for help from another member of staff
- recognising when emergency action is necessary
- administering salbutamol inhalers through a spacer
- keeping appropriate records of asthma attacks.

## ALL Members of Staff

In addition to this, it would be reasonable for **ALL** members of staff to be:

- trained to recognise the symptoms of an asthma attack and, ideally, how to distinguish them from other conditions with similar symptoms
- aware of this policy
- aware of how to check if a child is on the asthma register
- aware of how to access the emergency inhaler and who the designated members of staff are, and the policy on how to access their help

**Asthma UK has produced demonstration films on using a metered-dose inhaler and spacers suitable for staff and children.**

<http://www.asthma.org.uk/knowledge-bank-treatment-and-medicines-using-your-inhalers>

## What Arrangements are in Place at our School?

Medication is provided by the parents at the beginning of every school year or when needed. It is handed back at the end of the school year or on the expiration date whichever is sooner.

Children who have known allergies have a completed Health Plan. These plans are updated annually. All members of staff who work with that child are aware of the plan. Staff have regular training on the administration of medicines.

All medication for its treatment is kept unlocked in the classroom in a place which is easily accessible.

## **Healthcare Plan**

Pupils with asthma will need to have an individual healthcare plan, details about which are given on page 6 and in Appendix 3.

It is important to agree with parents of children with asthma how to recognise when their child's asthma gets worse and what action needs to be taken at that time. An Asthma Action Plan (available from Asthma UK) is a useful way to store written information about a child's asthma. The child's GP or Asthma Nurse will complete this in conjunction with the child and his/her parent. It includes details of the inhalers used, asthma triggers for the child, emergency action and contacts for the GP or Asthma Nurse. This can be attached to the healthcare plan.

In early years settings it is foreseeable that the younger children will not have the ability to convey to members of staff that their symptoms are getting worse or identify what medicines they need to take and when. It is, therefore, imperative that early years and primary school staff, who have younger children in their classes, know how to identify when symptoms of asthma are getting worse and what action they need to take when this happens. This should be reinforced by asthma action plans, asthma medicine cards (both provided by parents) and regular training and support for staff.

## **Asthma Register**

A register of children who have been diagnosed with asthma or prescribed a reliever inhaler should be kept in class. This is part of the medical needs register.

Schools should ensure that the asthma register is easy to access, and allows for a quick check to take place to establish if a child is recorded as having asthma and that consent for an emergency inhaler to be administered has been obtained.

## **Carrying the Medication**

Inhalers should be stored safely away and members of staff should issue them when the child needs the medication.

If the child is too young or immature to take personal responsibility for his/her inhaler, members of staff should make sure that it is stored in a safe but readily accessible place, and clearly marked with the child's name and photograph.

All asthma medicine should be clearly labelled with the child's name and photograph. The expiry date of the medicines should be checked every six months.

Where appropriate asthma inhalers should be carried by the young person at all times as often children recognise their own symptoms.

## **Emergency Salbutamol Inhalers in Schools**

As indicated above, schools are now permitted to keep a supply of salbutamol inhalers on site for use in an emergency. This is a sensible contingency arrangement in the event that children lose, forget or break their inhalers.

The emergency salbutamol inhaler should only be used by children:

- who have been diagnosed with asthma, and prescribed a reliever inhaler

- who have been prescribed a reliever inhaler
- for whom written parental consent for use of the emergency inhaler has been given.

Information on the use of the emergency inhaler should be recorded in a child's individual healthcare plan.

Keeping an inhaler for emergency use will have many benefits. It could prevent an unnecessary and traumatic trip to hospital and, potentially, save the child's life. Having a protocol that sets out how and when the inhaler should be used will also protect members of staff by ensuring they know what to do in the event of a child having an asthma attack; this should include:

- Establishing arrangements for the supply, storage, care and disposal of the inhaler and spacers. Assigning these responsibilities to at least two volunteers is recommended
- Maintaining a register of pupils who have been diagnosed with asthma or prescribed a reliever inhaler. The register should confirm that parental consent has been obtained for use of the emergency inhaler and a copy of it should be kept with the emergency inhaler
- Having written parental consent for use of the emergency inhaler included as part of a child's individual healthcare plan. This consent can either be secured by amending the School/Parental Agreement Form (Appendix 1) to include this permission or by using the specific consent form for use of the emergency inhaler (Appendix 6) which should be updated regularly, ideally annually, to take account of changes to a child's condition.
- Arranging for appropriate support and training for staff in the use of the emergency inhaler in line with this policy.
- Keeping a record of use of the inhaler (including when and where the attack took place, how much medication was given and by whom) and informing parents or carers that their child has used the emergency inhaler (this should be in writing so the parent can pass the information onto the child's GP – a sample letter is attached as Appendix 7)

The Medication Coordinator or Head should monitor the protocol to ensure compliance with it.

An emergency asthma inhaler kit should include:

- a salbutamol metered dose inhaler
- at least two single-use plastic spacers compatible with the inhaler
- instructions on using the inhaler and spacer
- instructions on cleaning and storing the inhaler
- manufacturer's information
- a checklist of inhalers, identified by their batch number and expiry date, with monthly checks recorded

- a note of the arrangements for replacing the inhaler and spacers
- a register of children permitted to use the emergency inhaler as detailed in their individual healthcare plans
- a record of when the inhaler has been used
- a copy of the school protocol on the use of the emergency salbutamol inhaler

### ***Salbutamol***

Salbutamol is a relatively safe medicine, particularly if inhaled, but all medicines can have some adverse effects. Those of inhaled salbutamol are well known, tend to be mild, temporary and not likely to cause serious harm. The child may feel a bit shaky or may tremble, or may say that they feel their heart is beating faster. The main risk of allowing schools to hold a salbutamol inhaler for emergency use is that it may be administered inappropriately to a breathless child who does not have asthma. It is essential, therefore, that schools follow the advice on page 14 in relation to whom the emergency inhaler can be used by.

***Children may be prescribed inhalers for their asthma which contain an alternative reliever medication to salbutamol (such as terbutaline). The salbutamol inhaler should still be used by these children if their own inhalers are not accessible – it will still help to relieve their asthma and could save a life.***

### ***Storage and Care of the Inhaler***

Schools should ensure that the inhaler and spacers are kept in a safe central location, such as the school office or staffroom, which is known to all members of staff, and to which they have access to at all times. However, the inhaler must be stored out of the reach and sight of children. The inhaler and spacer should not be locked away.

The inhaler should be stored at the appropriate temperature (in line with the manufacturer's guidelines), usually below 30<sup>o</sup>C, protected from direct sunlight and extremes of temperature. The inhaler and spacers should be kept separate from any individual child's inhaler; the emergency inhaler should be clearly labelled to avoid confusion with a child's inhaler. An inhaler should be primed when first used (for example, spray two puffs). As it can become blocked again when not used over a period of time, it should be regularly primed by spraying two puffs.

To avoid possible risk of cross-infection, the plastic spacer should not be reused. It can be given to the child to take home for future personal use. The inhaler itself, however, can usually be reused, provided it is cleaned after use. The inhaler canister should be removed, and the plastic inhaler housing and cap should be washed in warm running water, and left to dry in air in a clean, safe place. The canister should be returned to the housing when it is dry, and the cap replaced, and the inhaler returned to the designated storage place.

However, if there is any risk of contamination with blood (for example if the inhaler has been used without a spacer), it should also not be re-used but disposed of.

The two named volunteers should have responsibility for ensuring that:

- on a monthly basis the inhaler and spacers are present and in working order, and the inhaler has sufficient number of doses available
- replacement inhalers are obtained when expiry dates approach
- replacement spacers are available following use
- the plastic inhaler housing (which holds the canister) has been cleaned, dried and returned to storage following use, or replacements are available if necessary.

### ***Disposal***

The Council's Waste Management and Disposal Team has determined that owing to the very small quantities of emergency inhalers which Northumberland schools will produce, school will put them into its recycling and rubbish bins, as described below:

If schools have an NCC recycling collection then the empty metal canister can be recycled via these bins, as the Council does permit empty aerosols to be added to its collections

The plastic part of the inhaler should be placed into the rubbish bin, following which it will be sent with other waste to the 'energy from waste' plant

Schools requiring further information about this particular topic should call 0845 600 6400 and ask to speak to the Contracts and Commercial Team (Waste Management).

### **PE and Off-site Activities**

Children with asthma should participate in all aspects of school life, including physical activities. They need to take their reliever inhaler with them on all off-site activities and these should also be available during physical education and sports activities. Physical activity benefits children with asthma in the same way as other children. Swimming is particularly beneficial, although endurance work may need to be avoided. Some children may need to take their reliever asthma medicines before any physical exertion. Warm-up activities are essential before any sudden activity especially in cold weather. Particular care may be necessary in cold or wet weather.

The emergency inhaler kit should be easily accessible should the child's primary inhaler not be available.

### **Action During an Attack**

When a child has an attack they should be treated according to their individual healthcare plan or asthma card, as previously agreed. If the child does not have his/her prescribed reliever inhaler available, then the school's emergency inhaler can be used in the circumstances described previously. An ambulance should be called if:

- the symptoms do not improve sufficiently after 10 puffs on the inhaler
- the child is too breathless to speak
- the child is becoming exhausted
- 
-

the child has a blue/white tinge around the lips the child has collapsed

## Diabetes

---

### What is Diabetes?

Diabetes is a condition where the amount of glucose in the blood is too high because the body cannot use it properly.

### What Causes it?

Diabetes is a disorder caused when the pancreas produces an insufficient amount of the hormone insulin or when insulin production is absent. There are two main types of diabetes which are discussed briefly below:

### Type 1 Diabetes

Type 1 diabetes develops when the insulin-producing cells have been destroyed and the body is unable to generate any of the substance. It is treated with insulin either by injection or pump, a healthy diet and regular physical activity. The majority of affected children have Type 1 diabetes.

### Type 2 Diabetes

Type 2 diabetes develops when the body does not produce enough insulin or the insulin that is produced does not work properly.

This type of diabetes is treated with a healthy diet and regular physical activity, though medication (and/or insulin) is often required.

In both instances each child may experience different symptoms and these should be discussed when drawing up the healthcare plan.

## **What is the Treatment for the Condition?**

For most children diabetes is controlled by injections of insulin each day. Some children may require multiple injections, though it is unlikely that they will need to be given injections during school hours.

In some cases, the child's condition may be controlled by an insulin pump. Most children can manage their own injections, however, if doses are required at school then supervision may be required and a suitable, private place to inject will need to be identified.

It has become increasingly common for older children to be taught to count their carbohydrate intake and adjust their insulin accordingly. This means that they have a daily dose of long-acting insulin at home, usually at bedtime and then insulin with breakfast, lunch and evening meal, and before substantial snacks. The child is taught how much insulin to give with each meal, depending on the amount of carbohydrate eaten. The child is then responsible for administering injections and the regime to be followed would be detailed in the individual healthcare plan.

It is essential that children with diabetes make sure that their blood glucose levels remain stable. They may check their levels by taking a small sample of blood and using a small monitor at regular intervals. They may need to do this during the school lunch break, before PE or more regularly if their insulin needs to be adjusted. The majority of older children will be able to undertake this task without assistance and will simply need a suitable place to do it. However, younger children may need adult supervision to carry out the test and/or interpret the results.

When members of staff agree to administer blood glucose tests or insulin injections, they should be trained by an appropriate health professional, usually a specialist nurse with clinical responsibility for the treatment of the particular child.

## **What Arrangements are in Place at our School?**

Medication is provided by the parents at the beginning of every school year or when needed. It is handed back at the end of the school year or on the expiration date whichever is sooner.

Children who have known allergies have a completed Health Plan. These plans are updated annually. All members of staff who work with that child are aware of the plan. Staff have regular training on the administration of medicines.

All medication for its treatment is kept unlocked in the classroom in a place which is easily accessible.

Children with diabetes need to be allowed to eat regularly during the day. This may include eating snacks during class-time or prior to exercise.

Members of staff need to be made aware that if a child should miss a meal or snack he/she could experience a hypoglycaemic episode (commonly known as a 'hypo') during which the blood glucose level falls too low.

It is, therefore, important that staff should be aware of the need for children with diabetes to have glucose tablets or a sugary drink to hand. After strenuous activity a child may experience similar symptoms, in which case the teacher in charge of physical education or other sessions involving physical activity should be aware of the need to take appropriate action.

## What are the Signs of a Hypoglycaemic Episode?

Staff should be aware that the following symptoms, either individually or in combination, may be an indicator of low blood sugar:

- Hunger
- Sweating
- Drowsiness
- Pallor
- Glazed eyes
- Shaking or trembling
- Lack of concentration
- Irritability
- Headache
- Mood changes, especially angry or aggressive behaviour

Each child may experience different symptoms and this should be discussed when drawing up individual healthcare plans.

## Emergency Action

If a child experiences a 'hypo', it is very important that he/she is not left alone and that a fast acting sugar, such as glucose tablets, a glucose rich gel or a sugary drink is brought to the child and given immediately. Slower acting starchy food, such as a sandwich or two biscuits and a glass of milk, should be given once the child has recovered, some 10-15 minutes later.

An ambulance should be called if:

- The child's recovery takes longer than 10-15 minutes
- The child becomes unconscious

## Hyperglycaemia

Some children may experience hyperglycaemia, which is a high glucose level. The underlying cause of hyperglycaemia will usually be from loss of insulin producing cells in the pancreas or if the body develops resistance to insulin.

More immediate reasons for it include:

- Missing a dose of diabetic medication, tablets or insulin
- Eating more carbohydrates than the body and/or medication can manage
- Being mentally or emotionally stressed
- Contracting an infection

The symptoms of hyperglycaemia include thirst and the passing of large amounts of urine. Tiredness and weight loss may indicate poor diabetic control. If these symptoms are observed members of staff should draw these signs to the attention of parents. If the child is unwell, is vomiting or has diarrhoea this can lead to dehydration. If the child is giving off a smell of pear drops or acetone this may be a sign of ketosis and dehydration and he/she will require urgent medical attention.

Further information on this condition can be found on the [Diabetes UK](#) website.

## Epilepsy

---

### What is Epilepsy?

Epilepsy is characterised by a tendency for someone to experience recurrent seizures or a temporary alteration in one or more brain functions.

### What Causes it?

An epileptic seizure, sometimes called a fit, turn or blackout can happen to anyone at any time. Seizures can happen for many reasons and can result from a wide variety of disease or injury.

Triggers such as anxiety, stress, tiredness and illness may increase the likelihood that a child will have a seizure. Flashing or flickering lights and some geometric shapes or patterns can also trigger seizures. The latter is called photosensitivity and is very rare. Most children with epilepsy can use computers and watch television without any problem.

### What are the Signs of the Condition?

Seizures can take many different forms and a wide range of terms may be used to describe the particular seizure pattern that individual children experience.

What the child experiences depends on whether all of the brain is affected or the part of the organ that is involved in the seizure. Not all seizures involve loss of consciousness. When only a part of the brain is affected, a child will remain conscious with symptoms ranging from the twitching or jerking of a limb to experiencing strange tastes or sensations such as pins and needles. Where consciousness is affected; a child may appear confused, wander around and be unaware of their surroundings. They could also display unusual, such as plucking at clothes, fiddling with objects or making mumbling sounds and chewing movements. They may not respond if spoken to. Afterwards, they may have little or no memory of the seizure.

Most seizures last for a few seconds or minutes, and stop of their own accord. In some cases, seizures go on to affect all of the brain and the child loses consciousness. Such seizures might start with the child crying out, then the muscles becoming stiff and rigid. The child may fall down. Then there are jerking movements as muscles relax and tighten rhythmically. During a seizure breathing may become difficult and the child's colour may change to a pale blue or grey colour around the mouth. Some children may bite their tongue or cheek and may wet themselves.

After a seizure a child may feel tired, be confused, have a headache and need time to rest or sleep. Recovery times vary. Some children feel better after a few minutes while others may need to sleep for several hours.

Another type of seizure affecting all of the brain involves a loss of consciousness for a few seconds. A child may appear 'blank' or 'staring', and sometimes there will be fluttering of the eyelids. Such absence seizures can be so subtle that they may go unnoticed. They might be mistaken for daydreaming or not paying attention in class.

## **What is the Treatment for the Condition?**

The great majority of seizures can be controlled by anti-epileptic medication. It should not be necessary to take regular medicine during school hours.

## **What Arrangements are in Place at our School?**

Medication is provided by the parents at the beginning of every school year or when needed. It is handed back at the end of the school year or on the expiration date whichever is sooner.

Children who have epilepsy have a completed Health Plan. These plans are updated annually. All members of staff who work with that child are aware of the plan. Staff have regular training on the administration of medicines.

All medication for its treatment is kept in a red bum bag worn by a member of staff in the classroom.

An ambulance will be called as soon as possible.

## **Healthcare plan**

An individual healthcare plan is needed when a pupil has epilepsy.

Parents and health care professionals should provide information to the school's Medication Coordinator so that it can be incorporated into the individual healthcare plan, detailing the particular pattern of an individual child's epilepsy. If a child experiences a seizure whilst at school, details should be recorded and communicated to parents including:

- any factors which might possibly have acted as a trigger to the seizure – for example visual/auditory stimulation, anxiety or upset.
- any unusual 'feelings' which the child reported prior to the seizure
- the parts of the body demonstrating seizure activity, such as limbs or facial muscles
- the time when the seizure happened and its duration
- whether the child lost consciousness
- whether the child was incontinent

The above information will help parents to give the child's specialist more accurate information about seizures and their frequency. In addition, it should form an integral part of the school's emergency procedures and relate specifically to the child's individual healthcare plan. The healthcare plan should clearly identify the type or types of seizures, including descriptions of the seizure, possible triggers and whether emergency intervention may be required.

Children with epilepsy should be included in all activities. Extra care may be needed in some areas such as swimming or participating in science lessons. The Medication Coordinator should discuss any safety issues with the child and parents as part of the healthcare plan, and these concerns should be communicated to members of staff.

## **Emergency Action**

Information regarding emergency management is given in Appendices 9 and 10. Appendix 9 covers the procedures to be followed with regard to first aid for all seizures, whilst Appendix 10 covers procedures to be followed if the casualty is known to have epilepsy and has been prescribed buccal midazolam or rectal diazepam.

An ambulance should be called during a convulsive seizure if:

- it is the child's first seizure
- the child has injured him/herself badly
- the child has problems breathing after a seizure
- a seizure lasts longer than the period identified in the child's healthcare plan
- a seizure lasts for five minutes and members of staff do not know how long the seizures usually last for a particular child
- there are repeated seizures, unless this is usual for the child, as described in the child's health care plan

During a seizure it is important to make sure the child is in a safe position, not to restrict a child's movements and to allow the seizure to take its course. Putting something soft under the child's head during a convulsive seizure will help to protect it from injury.

Nothing should be placed in the child's mouth. After a convulsive seizure has stopped, the pupil should be placed in the recovery position and a member of staff should stay with him/her until the child has fully recovered.

## **Status Epilepticus**

Status epilepticus is a condition described as one continuous, unremitting seizure lasting longer than five minutes or recurrent seizures without regaining consciousness between them for greater than five minutes. It must always be considered a medical emergency.

A five minute seizure does not in itself constitute an episode of status and it may subsequently stop naturally without treatment. However, applying emergency precautions after the five minute mark has passed will ensure that prompt attention will be available if a seizure does continue. Such precautions are especially important if the child's medical history shows a previous episode of status epilepticus.

Any child not known to have had a previous seizure should receive medical assessment as soon as possible. Both medical staff and parents need to be informed of any events of this nature.

## **Emergency Medication**

Two types of emergency medication are prescribed to counteract status, namely:

- Rectal diazepam, which is given rectally (into the bottom). This is an effective emergency treatment for prolonged seizures.
- Buccal (oromucosal) midazolam. This is a new authorised treatment for prolonged acute convulsive seizures, which is placed via syringe into the buccal cavity (the side of the mouth between the cheek and the gum). It may be considered as an alternative to rectal diazepam for this purpose.

These drugs are sedatives which have a calming effect on the brain and are able to stop a seizure. In very rare cases, these emergency drugs can cause breathing difficulties so the person must be closely watched until they have fully recovered.

Training in the administration of buccal midolazam and rectal diazepam is essential and is provided by the specialist nurse with clinical responsibility for the treatment of the particular child. Special training should be updated annually.

## **Administration of Buccal Midazolam and Rectal Diazepam**

Any child requiring rectal buccal midolazam or diazepam should have his/her medication reviewed every year. As an additional safeguard, each child requiring buccal midolazam or rectal diazepam should have his/her own specific healthcare plan that will focus exclusively on this issue. All interested parties should be signatories to this document. An example is reproduced in Appendix 11 below.

Buccal midolazam and rectal diazepam can only be administered in an emergency if an accredited first-aider, trained in mouth to nose/mouth resuscitation, is easily accessible (that is only one or two minutes away). At least one other member of staff must be present as well.

Arrangements should be made for two adults to be present for such treatment, at least one of whom is the same sex as the child; this minimises the potential for accusations of abuse. The presence of two adults can also make it much easier to administer treatment. Staff should protect the dignity of the child as far as possible, even in emergencies.

Staying with the child afterwards is important as buccal midolazam and diazepam may cause drowsiness. Moreover, those who administer buccal midolazam and rectal diazepam should be aware that there could be a respiratory arrest. If breathing

does stop a shake and a sharp voice should usually start the child breathing again; if this does not work it will be necessary to give mouth to mouth resuscitation.

## Unacceptable Practice

---

Although school staff should use their discretion and judge each case on its merits whilst referencing the child's individual healthcare plan, it is not considered acceptable practice to:

- prevent children from easily accessing their inhalers and medication and administering their medication when and where necessary
- assume that every child with the same condition requires the same treatment
- ignore the views of the child or their parents; or ignore medical evidence or opinion (although this may be challenged)
- send children with medical conditions home frequently or prevent them from staying for normal school activities, including lunch, unless this is specified in their individual healthcare plans
- if the child becomes ill, send them to the school office or medical room unaccompanied or with someone unsuitable
- penalise children for their attendance record if their absences are related to their medical condition, such as hospital appointments
- prevent pupils from drinking, eating or taking toilet or other breaks whenever they need to in order to manage their medical condition effectively
- require parents, or otherwise make them feel obliged, to attend school to administer medication or provide medical support to their child, including assisting with toileting issues. No parent should have to give up working because the school is failing to support their child's medical needs
- prevent children from participating, or create unnecessary barriers which would hinder their participation in any aspect of school life, including school trips by, for example, requiring parents to accompany the child

## Complaints

---

Similarly to the stance adopted above, the DfE's statutory guidance requires that governing bodies ensure that the school's policy is crystal clear. It needs to set out how complaints concerning the support provided to pupils with medical conditions may be made and how they will be handled.

Should parents or pupils be dissatisfied with the support provided they should discuss their concerns directly with the school. If, for whatever reason, this does not

resolve the issue, they may make a formal complaint via the school's existing complaints procedure.

<http://www.blythhortongrange.co.uk/images/Policies/ComplaintsProcedure.pdf>

# **Administration of Medication to Pupils**

## **Agreement between Parents and School** (Appendix 1)

In order to keep the administration of medication to a minimum, the Head or Medication Coordinator should consider requesting that parents administer the daily doses out of school hours. However, if this is not possible it will be necessary for the school and parents to make a formal agreement to enable members of staff to administer medication to pupils during the school day by completing the form below.

In most cases only medication that the child's doctor has prescribed can be administered, hence school staff should not administer 'over-the-counter' medication. However, at the discretion of the Head, it is permissible for paracetamol to be administered provided that the practice is strictly controlled in the same way as is prescribed medication. Further information is given on page 6.

**Note: Medicines must be kept in the original container as dispensed by the pharmacy.**

### **Part 1 – To be Completed by Parent/Carer**

**To the Headteacher:**

(add name)

**School:**

My child (name) \_\_\_\_\_ Date of birth: \_\_\_\_\_

Class \_\_\_\_\_ has the following medical condition \_\_\_\_\_

I wish for him/her to have the following medicine administered by school staff, as indicated below:

Name of Medication:

Dose/Amount to be given:

Time(s) at which to be given:

Means of administration:

How long will the child require this medication to be administered?

Known side effects and any special precautions (please attach details)

Procedures to take in case of emergency (please attach details)

### **Emergency Contact 1**

Name: \_\_\_\_\_

Telephone \_\_\_\_\_

Work: \_\_\_\_\_

Home: \_\_\_\_\_

Mobile: \_\_\_\_\_

### **Emergency Contact 2**

Name: \_\_\_\_\_

Telephone \_\_\_\_\_

Work: \_\_\_\_\_

Home: \_\_\_\_\_

Mobile: \_\_\_\_\_

Relationship: \_\_\_\_\_

Relationship: \_\_\_\_\_

*I undertake to deliver the medicine personally to the Headteacher or Medication Coordinator and to replace it whenever necessary. I also undertake to inform the school immediately of any change of treatment that the doctor or hospital has prescribed.*

Name: \_\_\_\_\_ Signature: \_\_\_\_\_

Relationship to child: \_\_\_\_\_ Date: \_\_\_\_\_

### **Part 2 - To be completed by Headteacher/Medication Coordinator**

#### **Confirmation of agreement to administer medicine**

It is agreed that (*child*) \_\_\_\_\_ will receive (*quantity and name of medicine*)  
\_\_\_\_\_ every day at (*time medicine to be administered, for example, lunchtime or afternoon break*) \_\_\_\_\_.

(*Child*) \_\_\_\_\_ will be given medication or supervised whilst he/she takes it by  
(*name of member of staff*) \_\_\_\_\_.

This arrangement will continue until \_\_\_\_\_ (*either the end date for the course of medicine or until the parents instruct otherwise*).

Name: \_\_\_\_\_ Signature: \_\_\_\_\_  
Headteacher/Medication Coordinator

School: \_\_\_\_\_

### **Parental Request for Child to Carry and Self-administer Medicine**

(Appendix 2)

**This form must be completed by a parent/carer**

**To: Headteacher:**  
*(add name)*

**School:**  
*(add school name)*

Name of child: \_\_\_\_\_ Class: \_\_\_\_\_

Address: \_\_\_\_\_

Name of Medication: \_\_\_\_\_

Procedures to be taken in an emergency: \_\_\_\_\_

### **Contact Information**

*I would like my child to keep his/her medicine on him/her for use, as necessary.*

Name: \_\_\_\_\_ Signature: \_\_\_\_\_

Daytime Tel no(s): \_\_\_\_\_ Date: \_\_\_\_\_

Relationship to child: \_\_\_\_\_

**If more than one medicine is to be given a separate form  
should be completed for each one.**

# Healthcare Plan for a Pupil with Medical Needs

(Appendix 3)

<b>Details of Child and Condition</b>	
Name of child:	<i>Add photo here</i>
Date of birth:	
Class/Form:	
Medical Diagnosis/Condition:	
Triggers:	
Signs/Symptoms:	
Treatments:	
Has the Parental Consent Form been completed? <span style="float: right;">Yes/No</span> <i>(Medication cannot be administered without parental approval)</i>	
Date:	Review Date:
<b>Medication Needs of Child</b>	
Medication:	
Dose:	
Specify if any other treatments are required:	
Can the pupil self-manage his/her medication? Yes/No If Yes, specify the arrangements in place to monitor this:	
Indicate the level of support needed, including in emergencies: <i>(some children will be able to take responsibility for their own health needs)</i>	

Known side-effects of medication:

Storage requirements:

What facilities and equipment are required? (such as *changing table or hoist*)

What testing is needed? (such as *blood glucose levels*):

Is access to food and drink necessary? (where used to manage the condition): Yes/No Describe what food and drink needs to be accessed

Identify any dietary requirements:

Identify any environmental considerations (such as *crowded corridors, travel time between lessons*):

Action to be taken in an emergency (If one exists, attach an emergency healthcare plan prepared by the child's lead clinician):

#### **Other Requirements**

Detail any specific support for the pupil's educational, social and emotional needs

(for example, how absences will be managed; requirements for extra time to complete exams; use of rest periods; additional support in catching up with lessons or counselling sessions)

### Emergency Contacts

#### **Family Contact 1**

Name: \_\_\_\_\_

Telephone  
Work: \_\_\_\_\_

Home: \_\_\_\_\_

Mobile: \_\_\_\_\_

Relationship: \_\_\_\_\_

#### **Family Contact 1**

Name: \_\_\_\_\_

Telephone  
Work: \_\_\_\_\_

Home: \_\_\_\_\_

Mobile: \_\_\_\_\_

Relationship: \_\_\_\_\_

#### **Clinic or Hospital Contact**

Name: \_\_\_\_\_

Telephone:  
Work: \_\_\_\_\_

#### **GP**

Name: \_\_\_\_\_

Telephone:  
Work: \_\_\_\_\_

### Signatures

Signed

(Headteacher)

Signed

(Medication Coordinator)

## **Individual Staff Training Record – Administration of Medication** (Appendix 4)

Name: \_\_\_\_\_ Job: \_\_\_\_\_ School: \_\_\_\_\_

## **Record of Medication Administered in School (Appendix 5)**

**Name:** \_\_\_\_\_

**Parental Consent: Use of Emergency Salbutamol Inhaler Appendix 6**

School Name:		
Name of child:		
Date of birth:	Class/Form:	

**Child showing symptoms of asthma/having an asthma attack**

1. I can confirm that my child has been diagnosed with asthma/has been prescribed an inhaler [delete as appropriate].
2. My child has a working, in-date inhaler, clearly labelled with their name, which they will bring with them to school every day.
3. In the event of my child displaying symptoms of asthma, and if their inhaler is not available or is unusable, I consent for my child to receive salbutamol from an emergency inhaler held by the school for such emergencies.

Name: \_\_\_\_\_ Signature: \_\_\_\_\_

Date: \_\_\_\_\_ Relationship to child: \_\_\_\_\_

Address: \_\_\_\_\_

Daytime Tel no(s): \_\_\_\_\_

**Specimen letter to inform parents that the emergency salbutamol inhaler was used (Appendix 7)**

*(The Head or Medication Coordinator should feel free to tweak this letter so as to reflect the school's own tone and style when writing to parents)*

Child's name: \_\_\_\_\_

Class: \_\_\_\_\_

Date of Incident: \_\_\_\_\_

Dear [enter name of parent(s)]

I thought I would drop you a line to let you know that [enter child's first name] experienced problems with \*his/her breathing today. This happened when [enter details]

\*A member of staff helped [enter child's first name] to use \*his/her asthma inhaler.

\*Unfortunately, [enter child's first name] did not have \*his/her own asthma inhaler with \*him/her, so a member of staff helped \*him/her to use the school's emergency asthma inhaler, which contains salbutamol. [Enter child's first name] took [enter number] puffs on the inhaler.

\* Unfortunately, *[enter child's first name]* own asthma inhaler was not working, so a member of staff helped \*him/her to use the school's emergency asthma inhaler which contains salbutamol. *[Enter child's first name]* took *[enter number]* puffs on the inhaler.

Although *[enter child's first name]* soon felt a lot better, I think it might be a good idea if you were to take \*him/her to see the family doctor for a check-up.

Yours sincerely

*[Enter signature]*

\*Headteacher/Medication Coordinator

*[\*Delete as appropriate]*

# **Emergency Action in the Event of an Asthma Attack**

(Appendix 8)

- Keep calm and reassure the child
- Encourage the child to sit up and slightly forward
- Use the child's own inhaler – if not available, use the emergency inhaler
- Remain with the child while the inhaler and spacer are brought to them
- Immediately help the child to take two separate puffs of salbutamol via the spacer
- If there is no immediate improvement, continue to give two puffs at a time every two minutes, up to a maximum of 10 puffs
- Stay calm and reassure the child. Stay with the child until he/she feels better. The child can return to school activities when he/she feels better
- If the child does not feel better or you are worried at ANYTIME before you have reached 10 puffs, CALL 999 FOR AN AMBULANCE
- If an ambulance does not arrive in 10 minutes give another 10 puffs in the same way

# **Emergency Action: Epilepsy - First Aid for all Seizures**

(Appendix 9)

- Ensure that the child is out of harm's way. Move the child only if there is danger from sharp or hot objects or electrical appliances. Observe these simple rules and let the seizure run its course
- Check the time the child starts to fit
- Cushion the head with something soft (a folded jacket would do) but do not try to restrain convulsive movements
- Do not try to put anything at all between the teeth
- Do not give anything to drink
- Loosen tight clothing around the neck, remembering that this could frighten a semi-conscious child and should be done with care
- Arrange for other children to be escorted from the area, if possible
- Call for an ambulance if:
  - a seizure shows no sign of stopping after a few minutes
  - a series of seizures take place without the individual properly regaining consciousness
- As soon as possible, turn the child onto his/her side in the semi-prone (recovery/unconscious) position, to aid breathing and general recovery. Wipe away saliva from around the mouth
- Be reassuring and supportive during the confused period which often follows this type of seizure. If rest is required, arrangements should be made for this purpose
- If there has been incontinence cover the child with a blanket to prevent embarrassment. Arrange to keep spare clothes at school if this is a regular occurrence

## **If a child is known to have epilepsy:**

- It is not usually necessary for the child to be sent home following a seizure, but each child is different. If the Headteacher feels that the period of disorientation is prolonged, it might be wise to contact the parents. Ideally, a decision will be taken in consultation with the parents when the child's condition is first discussed, and a Healthcare Plan drawn up
- If the child is not known to have had a previous seizure medical attention should be sought
- If the child is known to have diabetes this seizure may be due to low blood sugar (a hypoglycaemic attack) in which case an ambulance should be summoned immediately

# **Emergency Action: First Aid for Children Known to Have Epilepsy and Prescribed Rectal Diazepam**

(Appendix)

- Ensure that the child is out of harm's way. Move the child only if there is danger from sharp or hot objects or electrical appliances. Observe these simple rules and let the seizure run its course.
- Check the time the child starts to fit
- Cushion the head with something soft (a folded jacket would do) but do not try to restrain convulsive movements
- Do not try to put anything at all between the teeth
- Do not give anything to drink
- Loosen tight clothing around the neck, remembering that this could frighten a semi-conscious child and should be done with care
- Arrange for other children to be escorted from the area, if possible
- Rectal diazepam must only be given to a child with a prescription that a Consultant Paediatrician has endorsed and updated annually
- Rectal diazepam must only be administered in an emergency by an appropriately trained member of staff in the presence of at least one other member of staff
- Rectal diazepam must only be administered if a trained First Aider is on site
- If the child has been convulsing for five minutes and there is no suggestion of the convulsion abating, the first dose of rectal diazepam should be given. The medication should indicate the name of child, the date of birth, date of expiry, contents and the dosage to be administered
- If after a further five minutes
  - (a) a seizure shows no sign of stopping or
  - (b) a series of seizures takes place without the individual properly regaining consciousness, then call an ambulance
- As soon as possible, turn the child onto his/her side in the semi-prone (recovery/unconscious) position to aid breathing and general recovery. Wipe away saliva from around the mouth
- Be reassuring and supportive during the confused period which often follows this type of seizure. Many children sleep afterwards and if rest is required, arrangements could be made for this purpose
- If there has been incontinence cover the child with a blanket to prevent embarrassment. Arrange to keep spare clothes at school if this is a regular occurrence
- A child should be taken home after a fit if he/she feels ill

# Individual Care Plan for the Administration of Rectal Diazepam

(Appendix)

This care plan should be completed by or in consultation with the medical practitioner (*Please use language appropriate to the lay person*)

## Details of Child and Condition

Name:

Class:

Date of birth:

Identify the seizure classification and/or description of seizures which may require rectal diazepam

*(Record all details of seizures, for example goes stiff, falls, convulses down both sides of body, convulsions last 3 minutes etc. Include information re: triggers, recovery time etc. If in status epileptics, note whether it is convulsive, partial or absence)*

Usual duration of seizure?

Other useful information:

## Diazepam Treatment Plan

**When should rectal diazepam be administered?** (*Note here should include whether it is after a certain length of time or number of seizures*)

**Initial dosage: how much rectal diazepam is given initially?** (*Note recommended number of milligrams for this person*)

**What are the usual reactions to rectal diazepam?**

**What action should be taken if there are difficulties in the administration of rectal diazepam such as constipation/diarrhoea?**

**Can a second dose of rectal diazepam be given? Yes/No**

If Yes, after how long can a second dose of rectal diazepam be given? (*state the time to have elapsed before re-administration takes place*)

How much rectal diazepam is given as a second dose? (*state the number of milligrams to be given and how many times this can be done after how long*)

**When should the person's usual doctor be consulted?**

**When should 999 be dialled for emergency help?**

- if the full prescribed dose of rectal diazepam fails to control the seizure Yes/No
- Other (Please give details)

**Who Should:**

- administer the rectal diazepam? (*ideally someone should be trained in at least 'Emergency Aid,' preferably 'First Aid at Work'*):
- witness the administration of rectal diazepam? (*this should normally be another member of staff of the same sex*):

**Who/where needs to be informed?**

Parent \_\_\_\_\_ Tel: \_\_\_\_\_

Prescribing Doctor:\_\_\_\_\_ Tel: \_\_\_\_\_

Other:\_\_\_\_\_ Tel: \_\_\_\_\_

**Precautions: under what circumstances should rectal diazepam not be used? (for example, Oral Diazepam already administered within the last.....minutes)**

***All occasions when rectal diazepam is administered must be recorded on the “Record of Use of Rectal Diazepam” log sheet (Appendix 12)***

**This plan has been agreed by the following:**

**Prescribing Doctor**

Name \_\_\_\_\_ Signature \_\_\_\_\_ Date \_\_\_\_\_

**Authorised person(s) trained to administer rectal diazepam**

Name \_\_\_\_\_ Signature \_\_\_\_\_ Date \_\_\_\_\_

Name \_\_\_\_\_ Signature \_\_\_\_\_ Date \_\_\_\_\_

**Parent**

Name \_\_\_\_\_ Signature \_\_\_\_\_ Date \_\_\_\_\_

**Headteacher/Medical Coordinator**

Name \_\_\_\_\_ Signature \_\_\_\_\_ Date \_\_\_\_\_

This form should be available at every medical review of the patient and copies held by the GP and the school.

**Expiry date of this form:** \_\_\_\_\_

**Copy holders to be notified of any changes by:** \_\_\_\_\_

# Record of Use of Rectal Diazepam (Appendix 12)

Name of Child: \_\_\_\_\_ Class: \_\_\_\_\_

Date:					
Recorded by:					
Type of seizure:					
Length and/or number of seizures:					
Initial dosage:					
Outcome:					
Second dosage (if any):					
Outcome:					
Observations:					
Parent informed:					
Prescribing doctor informed:					
Other information:					
Witness:					
Name of Parent re-supplying dosage:					
Date delivered to school:					